

# Metoclopramide

**BRAND NAME: REGLAN**

AVAILABLE IN  
5 mg & 10 mg  
TABLETS

(Oral suspension exists but is famous  
for unpalatable taste to pets)

## HOW THIS MEDICATION WORKS

Nausea can be the result of many medical scenarios including infection, poisoning, anxiety, vertigo, and primary stomach and intestinal disease. Recovery depends on proper nutrition and gastrointestinal function which in turn depend on the control of nausea. Obviously nausea control is important for general patient comfort as well.

Metoclopramide has two uses: nausea relief through its direct effect on the brain and improvement of the rhythm of stomach contraction (i.e. stomach motility modification.) In the case of metoclopramide, probably the anti-nausea effect is the dominating mechanism of action but it is important to understand the motility-modifying effect as well.

One of the stomach's most important functions is to grind the food we eat into a fine slurry that passes through the intestines freely. A strong rhythm of contraction is necessary to effect this and this rhythm creates the stomach's motility.

Motility disorders are common and may be chronic (of long duration) or of sudden onset. When motility is reduced in the stomach, food pools there and creates a sensation of nausea and bloating. In some cases, bile refluxes from the intestine back into the stomach causing irritation and more nausea. Metoclopramide normalizes stomach contractions so that food and bile can pass in the correct direction.

There is an additional effect of metoclopramide that is helpful in the control of nausea: There is a special biochemical barrier separating the brain and central nervous system from the blood stream so that only certain biochemicals can cross over. This partition is called the blood-brain barrier. Metoclopramide is able to cross the blood-brain barrier and act on the brain directly to control the sensation of nausea.

Metoclopramide thus helps the vomiting patient by normalizing stomach motility and acting directly on the brain to reduce the sensation of nausea.

## HOW THIS MEDICATION IS USED

Signs of nausea in pets include:

- Loss of appetite
- Vomiting
- Drooling
- Loud stomach or intestinal sounds
- Chewing motions

Metoclopramide can be used to control nausea in any situation (except in the event of stomach or intestinal obstruction). If a motility problem is suspected (a classical history would include vomiting food many hours after eating), metoclopramide should also be helpful. If a motility disorder extends beyond the stomach (for example, a colon motility disorder might lead to constipation), metoclopramide would not be expected to be helpful.

### **METOCLOPRAMIDE IS BEST GIVEN 20 MINUTES BEFORE A MEAL**

## **SIDE EFFECTS**

Side effects are rare with metoclopramide but generally stem from the ability to penetrate the blood-brain barrier. Animals predisposed to seizures (i.e., epileptics) may experience seizures though this should not be a problem for normal animals.

Some animals experience drowsiness and others will experience marked hyperactivity. Hyperactivity can be reversed with a dose of Benadryl (diphenhydramine).

Constipation has been reported as a side effect in some individuals. This is not a common side effect.

If you suspect your pet is having an adverse reaction to metoclopramide, discontinue the medication and inform your veterinarian. There is another motility modifier called cisapride (Propulsid) that may be a good alternative. This medication does not cross the blood-brain barrier and hence does not have the behavioral side effects that metoclopramide can, but it also does not reduce the sensation of nausea by this additional mechanism either.

## **INTERACTIONS WITH OTHER MEDICATIONS**

Metoclopramide should not be used with phenothiazine tranquilizers (such as acepromazine) if there is any potential for seizures in the patient. Both medications serve to lower the threshold to seizures.

Propantheline bromide (Pro-Banthine), which is used to treat some cardiac arrhythmias, will inactivate metoclopramide.

Narcotic analgesics may negate the effects of metoclopramide.

## **CAUTIONS AND CONTRAINDICATIONS**

In patients with pheochromocytoma (a rare adrenal tumor) the use of metoclopramide can induce dangerously high blood pressure.

If stomach obstruction is suspected in a given patient, it is best not to encourage motility and some other anti-nausea drug is preferred. Having said this, it is important to realize that as motility modifiers go, other medications are much stronger. If one is actually looking for motility modification as a chief effect and metoclopramide is felt to be inadequate, one of the following medications might be of use:

Cisapride      Erythromycin      Ranitidine      Tegaserod

Metoclopramide crosses the placenta and into milk and is probably best avoided in pregnant or nursing mothers.



## **Mar Vista Animal Medical Center**

3850 Grand View Blvd., Los Angeles, CA 90066 • (310) 391-6741 • Fax: (310) 391-6744  
Additional drug and general pet care information can be found on our world wide web site:  
<http://www.marvistavet.com>